


JPL PUBLICATION 82-44, VOLUME II

Planetary Geometry Handbook

Venus Positional Data, 1988-2020

Andrey B. Sergeevsky
Gerald C. Snyder
Barbara L. Paulson
Ross A. Cunniff



October 1, 1983

NASA

National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

The research described in this publication was carried out by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

Abstract

This document contains graphical data necessary for the analysis of planetary exploration missions to Venus. Positional and geometric information spanning the time period from 1988 through 2020 is presented; the text explains the data and its usage. This volume is one in a planned series covering planetary mission targets.

Preface

This publication is one in a series of volumes devoted to planetary positional and geometric data. The present Volume II provides information describing the planet Venus. Other volumes to be published in 1983 are III, IV, and V, describing Mars, Jupiter, and Saturn geometric data, respectively. The presentations of planetary positional and geometric data for Mercury (Volume I), Uranus (Volume VI), Neptune (Volume VII), and the Pluto/Charon System (Volume VIII) will be published later.

Contents

I. Introduction	1
II. Description of the Data	1
A. General Comments	1
B. Data Presentation	1
C. Definition of Terms	1
D. Description of Individual Plots	4
III. Application of the Data Presented	6
A. Geocentric Declination of Target Body	6
B. Planetocentric Declination of Earth and Sun	7
C. Right Ascension of Target Planet, Sun, and Earth	7
D. Sun- or Earth-to-Planet Distance	7
E. Heliocentric Longitude	7
F. Angles Between Sun, Earth, and Planet (SEP and ESP)	7
G. Cone and Clock Angles	8
H. Station Rise and Set	8
IV. Astrodynamic Constants	8
Acknowledgments	10
References	11
Figures	
1. Definition of vernal equinox	2
2. Definition of Sun- and Earth-to-planet distance, and Earth-Sun- and Sun-Earth-planet angles at any epoch	3
3. Definition of cone and clock angles	3
4. Schematic of a distant object's rise and set geometry, with a mask; polar view	4
5. Circumpolar and inaccessible regions of the sky for a station observer; meridional view	5
6. Definition of opposition and conjunction events involving Sun, Earth, and planet	8

7. Definition of the mean elements of a planetary orbit	9
8. IAU definition of planetary pole and prime meridian at time d_{50}	10

Tables

1. Deep Space Network tracking stations	5
2. Non-DSN tracking stations	5

Positional Data

Venus

1988–2020	13
1988	15
1989	21
1990	27
1991	33
1992	39
1993	45
1994	51
1995	57
1996	63
1997	69
1998	75
1999	81
2000	87
2001	93
2002	99
2003	105
2004	111
2005	117
2006	123
2007	129
2008	135
2009	141
2010	147
2011	153
2012	159

2013	165
2014	171
2015	177
2016	183
2017	189
2018	195
2019	201
2020	207

Positional Data

Earth

1985–2020	213
------------------	-----

I. Introduction

The purpose of this series of planetary geometry handbooks is to provide mission and science planners, as well as trajectory designers, with graphical information sufficient for preliminary mission design and evaluation. It is intended to be used in conjunction with the appropriate volumes and parts of the Mission Design Handbooks, Ref. 1, which describe trajectory aspects of the particular missions.

In most respects, this planetary geometry series represents a continuation of the second parts of each of the three volumes of Ref. 2. It extends their coverage by commencing in 1985 and continuing to the year 2020. This time span was chosen to provide sufficient mission duration flexibility for all Earth departures through 2005, as presented in Ref. 1.

The series will consist of eight volumes, each describing a planet as follows:

Volume	Target planet
I	Mercury
II	Venus
III	Mars
IV	Jupiter
V	Saturn
VI	Uranus
VII	Neptune
VIII	Pluto/Charon System

The present Volume II is devoted to data on the positions of the planet Venus. It presents information characterizing the geocentric and heliocentric positional geometry of a spacecraft while it is in the vicinity of the subject planetary body. The spacecraft could encompass a lander, an entry probe, or a flyby vehicle at encounter. Such wide usage of the data is possible because planetary positions change but slowly, while the distances to Earth or Sun are always large compared to those between the spacecraft and the target planet during the encounter.

II. Description of the Data

A. General Comments

All plots for this handbook series were computer generated, to reduce production cost. A program originally developed for Mariner 10 (MVM) and later used to produce the positional

plots for Ref. 2 has been used, with some modifications, on this series as well.

The major change in the new version of the program was the incorporation of the numerically integrated JPL Planetary DE-118 Ephemeris (Ref. 3) as the source for positional planetary information. An analytical and thus less-precise ephemeris, based on the mean elements of Ref. 4, was used in the preceding set of volumes, Ref. 2.

B. Data Presentation

The data in this volume are grouped into 36 subsections, each covering a specific calendar year. The data commence each year on January 1.0, 0^h Greenwich mean time (GMT) and continue through the year until "Day 380," thus providing a two-week overlap into the next year. There are eight plots presented for each year, labeled as follows:

- (1) *Declination* (DECLIN).
- (2) *Right ascension* (RT.ASC).
- (3) *Distance* (DISTANCE).
- (4) *Ecliptic longitude* (EC.LON).
- (5) *Sun-Earth-planet angles* (SUN-EARTH-PLANET).
- (6) *Cone and clock angles* (CA and KA, respectively).
- (7) *Station rise/set times* (STATION RISE/SET) for the three DSN stations (GMT, HRS).
- (8) Same as (7) for three other stations (Weilheim, Nobeyama, and Arecibo).

Twenty-six variables, relating positions of Earth, Sun, target planet (Venus), the star Canopus, and six tracking stations, are depicted on these plots for each subsection.

C. Definition of Terms

Planetary positional geometric data can be represented by directions to specified bodies and by angles between some of these directions, both shown as functions of calendar time. To distinguish among these bodies, the following subscripts are introduced:

- (1) E = Earth.
- (2) S = Sun.
- (3) P = planet (Venus in this volume).

Two fundamental planes are used in astrodynamics: that of the Earth's orbit about the Sun and that of the Earth's equator. The same two geometric concepts can be extended to all celestial bodies.

The plane of the mean Earth's orbit (EMO) is also called the ecliptic; the term "mean" refers to an averaging process, which eliminates minor short-period oscillations of the orbit pole; these oscillations are caused by repetitive perturbing accelerations caused by solar system objects other than the Sun.

The rotational, angular-momentum vector of each planet defines the polar-axis orientation of that celestial body, but not necessarily its north pole. The plane normal to this axis of rotation is defined as the planetary equator of that body. The latest (1983) International Astronomical Union (IAU) definition of planetary poles is such that the *north pole* of a body is that pole that points north of the Invariable (or Laplacian) Plane, defined as the plane normal to the total angular momentum vector of the solar system. This plane is very close to the ecliptic, and deviates from the latter by less than 0.5 deg (Ref. 5).

The point on the celestial sphere where the ecliptic crosses the Earth's mean equator (EME) from south to north is called the *ascending node* of the Ecliptic, while the line passing from the Earth's center through that point is called *Earth's vernal equinox* (Υ). The mean Sun is positioned at that point of the sky (as seen from Earth's center) at the time of each spring equinox.

Both the equator and the orbit plane of each planet slowly precess on their respective reference planes, driven by "secular" perturbing effects due to other solar system bodies.

The mean ecliptic regresses, i.e., its nodes slide backwards, on the Invariable Plane at the very slow rate of about 47 seconds of arc per century, while the Earth mean equator regresses on the ecliptic at about 50.26 seconds of arc per annum.


Clearly these motions affect the position of vernal equinox, which is therefore specified as being "of" an epoch, e.g., "of date" (i.e., the present instant of time), or "of 1950.0." The Besselian (tropical) new year has for a long while been considered a convenient defining epoch, usually advanced each half-century, e.g., 1950.0, recently renamed B1950.0, which occurred on January 0.923357, 1950 (i.e., slightly before civil new year 1950, celebrated at 0^h GMT, on January 1.0). The acronym EME50 therefore refers to the Earth mean equator and equinox of B1950.0, while EMO50 specifies the ecliptic plane of that epoch.

Rather than refer to dates frequently quoted in differing calendars, it is customary to specify dates in Julian Days (JD), consecutively counted from an arbitrary date in antiquity (January 1.0, 4713 B.C.). Thus the epoch JD 243 3282.423357 is equivalent to B1950.0.

For each celestial body, a *vernal equinox* may be similarly defined as the ascending node of that body's mean orbit plane (about its primary) upon the body's equator plane. This preferred direction is usually taken to be the X-axis of the inertial coordinate system, centered on that body. The Z-axis coincides with the north-pole axis of the body, and the Y-axis completes a *right-handed* coordinate system.

The equations and related constants defining the above planes are presented in Section IV and may be found in Refs. 4 and 5. Data in Ref. 5 reflect the latest IAU definitions.

A useful way to describe celestial directions is by means of a spherical coordinate system, based either on the equator or else on the mean orbit plane of a body, e.g., that of the Earth or of the target planet. Two angles, declination and right ascension, define the position of any point on the celestial sphere, referred to a planetary equator (Fig. 1).



$$\vec{R}(x, y, z) = (\cos \delta \cdot \cos \alpha, \cos \delta \cdot \sin \alpha, \sin \delta)$$

Fig. 1. Definition of vernal equinox; a position vector \vec{R} , its declination δ , and right ascension α

Declination (δ) is the central angle formed by the object's position vector with the equator plane, i.e., it is an elevation angle with respect to that plane. It is a form of latitude measure, and is positive towards the north pole and negative to the south of the equator.

Right ascension (α) of an object is the angle in the equator plane between vernal equinox and the projection of the

object's position vector into the equator plane. It is positive in the eastward (i.e., right-handed, counterclockwise, when seen from north) direction, regardless of the planet's (celestial body's) own sense of rotation.

Both of these angles are expressed on the unit sphere. Actual distances between Sun, Earth, and planet can be presented separately and are measured in astronomical units (AU) - a defined mean distance between Sun and Earth (1 AU = 149,597,871 km; see Astrodynamic Constants, Section IV) - or also in kilometers. The distance plot scale in this volume of the handbook is in units of millions of kilometers ($10^6 \times \text{km}$).

Position angles on the celestial sphere expressed with respect to a planetary orbit plane (e.g., the ecliptic) are referred to as latitudes and longitudes.

Ecliptic longitude (L) of an object is the angular distance between Earth's vernal equinox and the projection of the object's position vector into the ecliptic (EME50), measured in an eastward (i.e., counterclockwise) positive direction.

Sun-Earth-planet (SEP) angle and the corresponding *Earth-planet-Sun (EPS)* angle are defined by the two respective vector pairs: the Earth-Sun and Earth-planet vectors for the SEP-angle, and the Sun-Earth and Sun-planet vectors for the EPS-angle (Fig 2). The angles are important aids in avoiding periods of communication blackout.




Fig. 2. Definition of Sun- and Earth-to-planet distance, and Earth-Sun- and Sun-Earth-planet angles at any epoch

Cone and clock angles (CA and KA, respectively) are spacecraft-centered coordinates of an object. The polar Z-

axis of the system corresponds to the longitudinal axis of the spacecraft. It is usually pointed either towards the Sun (solar panels must generally be normal to the Sun direction), or to the Earth (the parabolic antenna bore axis should point along the communication link, i.e., towards the Earth). Within this book, the planet-Sun direction is taken to be the Z-axis, orientation of the CA, KA system (Fig. 3).




Fig. 3. Definition of cone and clock angles

The prime meridian of this system contains the viewing slot of the spacecraft star tracker, which is usually, but not necessarily, oriented toward the bright southern star Canopus (α -Carinae). Its position on January 1.0, 1950, was at right ascension $\alpha = 95.7103875$ and declination $\delta = -52.667625$ deg, expressed with respect to the EME50. Stellar proper motions slowly change star locations; the method for updating stellar positions as well as data for other stars are provided in the FK4 Star Catalogue (Ref. 6).

ORIGINAL PAGE IS
OF POOR QUALITY

The cone angle of an object, described by its spacecraft-centered position vector, is its angular distance from the Z-axis; i.e., it is the Sun-planet-object angle.

The clock angle of an object is measured in the X-Y plane, which is akin to a "spacecraft equator" and normal to the spacecraft Z-axis. Clock angles are measured from the star-tracker (Canopus) meridian; they are positive clockwise when looking along the Z-axis towards the Sun.

Although the clock angle of Canopus is zero by definition (when the spacecraft is Canopus oriented), the Canopus cone angle generally differs from 90 deg, because the position of Canopus is offset several degrees from the ecliptic pole.

Station rise/set times are determined by the daily rotation of the Earth, which takes all ground tracking stations around the polar axis on the small circles that correspond to their terrestrial latitudes. As a result, every object on the celestial sphere rises and sets upon the local horizon of the station (Fig. 4), provided the object's declination is within ± 90 deg of the station latitude (e.g., an object at -55 deg declination cannot be seen by a 40 -deg latitude station). Only objects whose declinations equal the station latitude rise to the zenith, in this daily spectacle (Fig. 5). The angular height of an object above the local horizon is termed *elevation* (Γ). Within this book, an object is considered to have risen if it

reaches above an elevation mask of 6 deg, which accounts for obstructions by local terrain features, atmospheric signal refraction, and radio noise.

Rise and set time information is presented for the Deep Space Network (DSN), the stations for which are listed in Table 1, as well as for three other stations (Table 2).

The Arecibo radio telescope, located within a natural bowl, is not movable. The slewing of a dish, therefore, is replaced by horizontal motion of a cable-mounted cab, which translates the antenna feed to track a celestial object. The resulting daily coverage of this station is clearly very limited.

D. Description of Individual Plots

The individual plots within each annual subsection present the following 26 planetary geometric characteristics as functions of calendar time (recall that for the purposes of this volume "planet" refers to Venus):

Plot 1. DECLIN. This plot shows three curves:

- (1) The geocentric declination of the target planet, referred to the Earth mean equator and equinox of B1950.0 (EME50); it is labeled "P". All angular measures on the plots are in degrees.




Fig. 4 Schematic of a distant object's rise and set geometry, with a mask; polar view



Name	Geocentric latitude ϕ_g	Geocentric longitude (west) λ_g
DSS 14, Goldstone, California	35.24	116.89
DSS 43, Canberra, Australia	-35.22	211.02
DSS 63, Madrid, Spain	40.24	4.25

Name	Geocentric latitude ϕ_s	Geocentric longitude (west) λ_s
Weilheim, Germany	47 69	348 92
Nobeyama, Japan	35 94	221 53
Arecibo, Puerto Rico	18 34	66 75

- (7) The Earth-planet distance, in this volume expressed in units of $10^6 \times \text{km}$ (millions of km). It is labeled "EP".

- (8) The Sun-planet distance, also in millions of km. It is labeled "SP".

Plot IV. LC.LON: A single curve is shown on this plot:

- (9) The planet's heliocentric, ecliptic longitude (with respect to the mean ecliptic and equinox of B1950.0, i.e., EMO50). It is labeled "P".
The annual variation in Earth's heliocentric ecliptic longitude is repetitive to a fair degree of accuracy, hence only a single plot, valid throughout the 1985 to 2020 time period, is provided at the end of the plotted positional data section.

Plot V. SUN-EARTH-PLANET: Two curves are presented.

- (10) The "SEP" angle (Sun-Earth-planet)
(11) The "ESP" angle (Earth-Sun-planet)

Plot VI. CA, KA: Three curves are depicted on the plot:

- (12) The cone angle of Earth for a Sun-oriented spacecraft, labeled "ECA".
(13) The clock angle of Earth, for a Sun-Canopus two-axis-oriented spacecraft. It is labeled "EKA".
(14) The cone angle of Canopus, in any Sun-oriented spacecraft, attitude. It is labeled "CCA".

Plot VII. STATION RISE/SET: This plot shows six curves, each representing the Greenwich mean time (GMT) of the following daily Deep Space Network events (all based on a 6-deg mask above the local horizon).

- (15) Planet rise at DSS 14 (Goldstone), labeled "RISE 14".
(16) Planet set at DSS 14, labeled "SET 14".
(17) Planet rise at DSS 43 (Canberra), labeled "RISE 43".
(18) Planet set at DSS 43, labeled "SET 43".
(19) Planet rise at DSS 63 (Madrid), labeled "RISE 63".
(20) Planet set at DSS 63, labeled "SET 63".

Plot VIII. STATION RISE/SET: The plot presents up to six curves, representing the GMT of the following tracking station events (6-deg mask, unless specified otherwise):

- (21) Planet rise at Weilheim, Germany. It is labeled "R WEIL".
(22) Planet set at Weilheim, labeled "S WEIL".
(23) Planet rise at Nobeyama, Japan. It is labeled "R JAPN".

- (24) Planet set at Nobeyama, labeled "S JAPN".

- (25) Planet rise at the Arecibo radio telescope, Puerto Rico, above the 70-deg special horizon mask, if realizable. It is labeled "R AREC".

- (26) Planet set at Arecibo, under the same conditions as those in (25). It is labeled "S AREC".

III. Application of the Data Presented

The number of practical mission analysis problems to which the information presented in this volume is applicable is obviously large. By no means is it intended to enumerate herein a complete list of such applications. Rather, only some examples, typical of such usage, shall be discussed.

The comments will be arranged in the same order as the graphical output data are presented. Problems in mission analysis to which a particular output type is relevant will be discussed accordingly.

A. Geocentric Declination of Target Body

The spacecraft orbit determination process uses DSN radiometric range and range-rate (Doppler) data. Whenever the observed body as seen from Earth is at near-equatorial (≈ 0 deg) declination, a situation develops that seriously degrades orbit determination accuracy. As a consequence, the timing of critical maneuvers or planetary encounter events should either utilize more sophisticated data types (e.g., Δ DOR, a method based on long-base interferometry) or avoid periods when the target planet is near its node with the Earth's equator (i.e., the times of near-zero declination of the target body).

Communication with the spacecraft greatly depends on a ground station's ability to receive the data signal. An elevation angle high over the horizon, coupled with as long a listening period as possible, are two important assets in this respect. The highest elevation angle of an object at each station occurs at the meridian passage of the object; its value is:

$$\Gamma_{p \max} = 90 - |\phi_s - \delta_p|$$

Its maximum of $\Gamma_p = 90$ deg can be realized only at a station whose latitude, ϕ_s , equals the declination δ_p of the target. Exact equality, however, may cause some data loss: to continue tracking, some antennas may have to swing through a 180-deg arc in azimuth as soon as the object reaches zenith (i.e., at elevation $\Gamma_p = 90$ deg). As already mentioned, the maximum elevation angle drops with an increase in the differ-

ence between station latitude, ϕ_s , and object declination δ_p . At a difference of

$$|\phi_s - \delta_p| + \text{mask} \geq 90 \text{ deg}$$

coverage ceases altogether (Fig. 5). The mask, it will be recalled, is an angle of about 6 deg, designed to assure good signal strength at low elevations.

A notable aspect of this is that, as a station moves in latitude toward either pole (north or south), objects located between the nearer pole and the circumpolar latitude circle δ_c of the station, defined by

$$|\delta_c| = 90 - |\phi_s|$$

will never set and communication may be continuous, if so desired. Knowledge of target-planet declination allows one to choose the particular station for which most critical events would appear at high elevation angles or, conversely, to locate latitudes at which ships and portable or permanent stations should be positioned to support a given difficult mission tracking assignment.

B. Planetocentric Declination of Earth and Sun

These data are based on the target planet's own equator and provide a means of determining the solar lighting conditions at a landing site or during approach. They also allow one to obtain the Earth-oriented downlink elevation angle for a lander on the surface at a given latitude.

For ringed planets (e.g., Saturn), the data can indicate which side (north or south) of the rings is illuminated, which side is visible from Earth, and where on the planetary disk the ring will project its shadow. Observability of disk features and phenomena (e.g., the spokes) may also depend on this information.

C. Right Ascension of Target Planet, Sun, and Earth

This information, when combined with the declination data already discussed, provides positional reference directions toward these bodies, in spherical coordinates as presented, or in Cartesian unit vector form, if transformed as follows:

$$X = \cos \delta_0 \cos \alpha_0$$

$$Y = \cos \delta_0 \sin \alpha_0$$

$$Z = \sin \delta_0$$

where α_0 = right ascension and δ_0 = declination of vector to object.

A number of mission-related problems in spherical trigonometry may be solved using the above information.

D. Sun- or Earth-to-Planet Distance

When the spacecraft is near the planet, Sun-to-planet distance data help in computing such solar effects upon the spacecraft as solar heating and thermal balance, solar-panel electrical output, and spacecraft trajectory and attitude perturbation magnitudes due to the solar-radiation pressure.

Earth-to-planet distance information allows estimation of maximum available data transmission rates in the radio up- or downlink, as well as the one- and two-way light times required in communicating with the spacecraft.

When combined with information on direction unit vectors (X, Y, Z) from paragraph C above, position vectors for Sun, Earth, and planet may be constructed in the respective coordinate systems.

E. Heliocentric Longitude

The heliocentric longitudes of the departure and arrival planetary positions allow sketching the transfer trajectory trace and its orientation with respect to the ecliptic nodes or the perihelion of the target-planet orbit. The relevant planetary information, e.g., the planetary mean orbital elements and related constants, may be found in Section IV.

F. Angles Between Sun, Earth, and Planet (SEP and ESP)

These angles are of paramount importance to spacecraft communication planning. For example, the radio downlink signal becomes distorted and contaminated with noise, if the radio beam passes through the solar corona. This effect is observed near "superior conjunction" between Sun and the target planet. It occurs when the Sun-Earth-planet angle (SEP) reaches near-zero values (commonly, $\leq \pm 5$ deg) while at the same time the Earth-Sun-planet angle (ESP) reaches its maximum value near 180 deg. This unfavorable condition as well as the two other extreme configurations of "opposition" and "inferior conjunction" (neither of which significantly affects up- and downlink communication capability) are shown in Fig. 6.

It is essential to proper mission design that important science, orbit determination, or engineering events be avoided

ORIGINAL PAGE IS
OF POOR QUALITY




Fig. 6. Definition of opposition and conjunction events involving Sun, Earth, and planet

during the 10 to 20 days that the spacecraft and/or target planet are near superior conjunction. On the other hand, solar physicists do delight in studying the corona's signature superimposed on the distorted signal. This period of time then becomes an opportunity for corona radio-occultation experiments and related solar studies, including relativity investigations.

G. Cone and Clock Angles

As outlined earlier, the cone- and clock-angle system is primarily useful because it is a spacecraft-fixed set of coordinates. As such, and together with data and equations provided in the Mission Design Handbook series (Ref. 1), the cone and clock angles presented on the plots can be used to resolve a number of geometry problems, such as:

- (1) The angular travel range required by the movable antenna to track the Earth, if Sun-oriented.
- (2) The extent of the Sun-sensor bias required, if the Earth-orientation of the entire spacecraft, together with its antenna, is desired.
- (3) The impact of lighting and thermal constraints upon spacecraft orientation options.

- (4) The Star tracker slot size in cone angle (CA) required to observe Canopus at different times during an orbiter mission.

H. Station Rise and Set

Station rise and set times, given in GMT, greatly affect mission design planning. After consideration is given to the one-way light time required by the radio signal to traverse the Earth-to-planet distance (and back if needed), significant spacecraft events can be positioned in universal (GMT) time using the rise/set information provided, such as to assure that a given station is "up," i.e., that the spacecraft is above mask in elevation angle at a suitable tracking station. The width (in hours) of the up-time band of a station is in itself an indication of the maximum elevation angle available during the station's pass "under" the spacecraft (i.e., the planet) — the greater the width, the higher the maximum elevation angle.

IV. Astrodynamic Constants


This section presents information necessary for relating the various coordinate systems used in the handbook. For other astrodynamic constants, see Refs. 1, 4, 5, and 7.

Mean orbital elements for the planets are *averaged* analytical models of planetary motion, adopted for reference usage by the astronomical community (e.g., H. M. Nautical Almanac Office or the U.S. Naval Observatory). Mean elements change as functions of time, T_{50} .

T_{50} is defined as the time interval, in Julian centuries, elapsed at date of interest, since epoch January 1.0, 1950, 0^h ephemeris time (ET) (i.e., Julian epoch J1950.0 = JD2433282.5). A Julian century contains 36525.0 mean solar (i.e., calendar) days. Ephemeris time is an atomically controlled uniform time, approximated as: ET = GMT + DUT, where the time increment DUT = +54.2 s (as of July 1983). DUT is currently increasing at about one second per year.

Six elements completely define the mean orbit (Fig. 7), as follows (all angular measures in degrees):

(a) CELESTIAL SPHERE VIEW



(b) ORBIT PLANE VIEW




Fig. 7. Definition of the mean elements of a planetary orbit

a = semimajor axis (in astronomical units, 1 AU = 149,597,871 km).

e = eccentricity of orbit (dimensionless).

M = mean anomaly, a mean motion angular position from perihelion (i.e., closest approach to Sun) at time T_{50} (for numerical convenience also expressed as d_{50} , days from January 1.0, 1950 0^h ET). True anomaly (actual position angle from perihelion) can be obtained from M by way of iteration using Kepler's transcendental equation.

i_{50} = inclination angle of mean orbit plane at time T_{50} , with respect to the mean ecliptic of epoch B1950.0 (EMO50).

Ω_{50} = longitude of ascending node of mean orbit upon the ecliptic of B1950.0, measured from Earth's vernal equinox.

ω_{50} = argument of perihelion, an angle measured from the ascending node of the mean orbit on the ecliptic of B1950.0 to the perihelion of the mean orbit of the planet.

As presented by F. Sturms (Ref. 4) and based on information in the Explanatory Supplement to the Ephemeris (Ref. 7), the mean elements are:

(1) Earth:

$$a = 1.00000023 \text{ AU} = 149,597,905 \text{ km}$$

$$e = 0.0167301085 - 0.000041926 T_{50} - 0.000000126 T_{50}^2$$

$$M = 358.000682 + 0.9856002628 d_{50} - 0.0001550000 T_{50}^2 - 0.0000033333 T_{50}^3$$

$$i_{50} = 0.013076 T_{50} - 0.000009 T_{50}^2$$

$$\Omega_{50} = 174.40956 - 0.24166 T_{50} + 0.00006 T_{50}^2$$

$$\omega_{50} = 287.67097 + 0.56494 T_{50} + 0.00009 T_{50}^2$$

(2) Venus:

$$a = 0.7233316 \text{ AU} = 108,208,867 \text{ km}$$

$$e = 0.00679684275 - 0.000047649 T_{50} + 0.000000091 T_{50}^2$$

$$M = 311.505478 + 1.6021301892 d_{50} + 0.0012860555 T_{50}^2$$

$$i_{50} = 3.39413 - 0.00086 T_{50} - 0.00003 T_{50}^2$$

$$\Omega_{50} = 76.22967 - 0.27785 T_{50} - 0.00014 T_{50}^2$$

$$\omega_{50} = 54.63793 + 0.28818 T_{50} - 0.00115 T_{50}^2$$

The equatorial plane of a celestial body at an epoch of interest, T_{50} , was defined by the International Astronomical Union Working Group on Cartographic Coordinates and Rotational Elements (Ref. 5) by the right ascension α_{50} and declination δ_{50} of the body's "north" polar axis direction with respect to the Earth mean equator of epoch B1950.0 (EME50), as shown in Fig. 8. The position of the rotating prime meridian of the body is defined by an angle W , measured positive counterclockwise in that body's equatorial plane from the ascend-




Fig. 8. IAU definition of planetary pole and prime meridian at time d_{50} , elapsed since January 1.0, 1950 0^h ET

ing node of that equator upon the EME50, to the prime meridian position at epoch d_{50} . According to the Ref. 5, Table 1.

(1) Sun pole.

$$\alpha_{50} = 285.90 \text{ (deg)}$$

$$\delta_{50} = 63.90$$

$$W = 240.90 + 14.1844000 d_{50}$$

(2) Earth pole:

$$\alpha_{50} = 0.00 - 0.640 T_{50}$$

$$\delta_{50} = 90.00 - 0.557 T_{50}$$

$$W = 99.87 + 360.9856123 d_{50}$$

(3) Venus pole:

$$\alpha_{50} = 272.80$$

$$\delta_{50} = 67.20$$

$$W = 213.63 - 1.4814205 d_{50} \text{ (retrograde)}$$

(4) Invariable Plane pole:

$$\alpha_{50} = 272.40$$

$$\delta_{50} = +66.99$$

Note that in the computation of W , care should be taken not to lose significant digits during arithmetic calculations.

Positional data in this handbook were obtained using the JPL DE-118 precision planetary ephemeris on magnetic tape in tabular format (Ref. 3).

Acknowledgments

The contributions, reviews, and suggestions by members of the Handbook Advisory Committee, especially those of K. T. Nock, R. E. Diehl, W. I. McLaughlin, W. E. Bollman, P. A. Penzo, R. A. Wallace, D. F. Bender, D. V. Byrnes, L. A. D'Amario, T. H. Sweetser, and R. S. Schlaifer are acknowledged and greatly appreciated. The authors would like to thank Mary Fran Buehler and David E. Fulton for their editorial contribution.

References

1. Sergeyevsky, A. B., et al., *Interplanetary Mission Design Handbook*, JPL Publication 82-43, Volume 1, Parts 1-4, Jet Propulsion Laboratory, Pasadena, Calif., March 1, 1983.
2. Sergeyevsky, A. B., *Mission Design Data for Venus, Mars, and Jupiter through 1990*, Technical Memorandum 33-736, Jet Propulsion Laboratory, Pasadena, Calif., Sept. 1, 1975.
3. Newhall, XX, Standish, E. M., and Williams, J. G., "DE-102: A Numerically Integrated Ephemeris of the Moon and Planets, Spanning 44 Centuries," *Astronomy and Astrophysics*, 1983 (in press).
4. Sturms, F. M., *Polynomial Expressions for Planetary Equators and Orbit Elements With Respect to the Mean 1950.0 Coordinate System*, Technical Report 32-1508, pp. 6-9, Jet Propulsion Laboratory, Pasadena, Calif., Jan. 15, 1971.
5. Davies, M. E., et al., "Report of the IAU Working Group on Cartographic Coordinates and Rotational Elements of the Planets and Satellites: 1982," *Celestial Mechanics*, vol. 29, No. 4, pp. 309-321, April 1983.
6. Fricke, W., et al., *Fourth Fundamental Catalogue (FK 4)*, No. 10 in the series, *Veröffentlichungen des Astronomischen Rechen-Instituts Heidelberg*, Verlag L. Braun, Karlsruhe, 1963.
7. "Explanatory Supplement to the Ephemeris", Her Majesty's Stationery Office, London, 1961.

Positional Data

**Venus
1988--2020**

PRECEDING PAGE BLANK NOT FILMED

PAGE 12 INTENTIONALLY BLANK

Venus

1988

PRECEDING PAGE BLANK NOT FILMED



PAGE 14 INTENTIONALLY BLANK

DECLIN RT.ASC 1988

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

1988





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

1988

DISTANCE
EC.LON
1988





SEP, ESP
CA, KA
1988

VENUS

1988


ORIGINAL PAGE
OF POOR QUALITY



STAR/S
DSN
1988

ORIGINAL DATA SET
OF POOR QUALITY


VENUS 1988



**STA R/S
NON-DSN
1988**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1988



Venus



1989

DECLIN RT.ASC 1989

VENUS

1989

ORIGINAL PAGE 17
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

1989

**DISTANCE
EC.LON
1989**





SEP, ESP
CA, KA
1989

VENUS

1989


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1989

ORIGINAL PAGE 15
OF POOR QUALITY


VENUS 1989



STA R/S
NON-DSN
1989

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1989



Venus



1990

DECLIN RT.ASC 1990

VENUS

1990

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

1990

**DISTANCE
EC.LON
1990**





SEP, ESP
CA, KA
1990

VENUS

1990


ORIGINAL PAGE
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STA R/S
DSN
1990


VENUS 1990



**STAR/S
NON-DSN
1990**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1990



Venus



1991

DECLIN RT.ASC 1991

VENUS

1991

ORIGINAL PARTIAL
OF POOR QUALITY





ORIGINAL SET 103
OF FOUR QUEST?

VENUS

1991

**DISTANCE
EC LON
1991**





SEP, ESP
CA, KA
1991

VENUS

1991


ORIGINAL PAGE 10
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STAR/S
DSN
1991


VENUS 1991



STA R/S
NON-DSN
1991

ORIGINAL PAGE 1
OF POOR QUALITY

VENUS 1991



Venus



1992

DECLIN RT.ASC 1992

VENUS

1992

ORIGINAL FIGURE
OF POOR QUALITY





ORIGINAL FACSIMILE
OF POOR QUALITY

VENUS

1992

**DISTANCE
EC.LON
1992**





SEP, ESP
CA, KA
1992

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS


1992



ORIGINAL PAGE IS
OF POOR QUALITY

STAR/S
DSN
1992


VENUS 1992



STA R/S
NON-DSN
1992

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1992



Venus



1993

DECLIN RT.ASC 1993

VENUS

1993

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

1993

DISTANCE
EC.LON
1993





SEP, ESP
CA, KA
1993

VENUS

1993


OF POOR QUALITY



STAR/S
DSN
1993

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1993




**STA R/S
NON-DSN
1993**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

1993



Venus



1994

DECLIN RT.ASC 1994

VENUS



1994

ORIGINAL PAGE IS
OF POOR QUALITY



VENUS

1994



ORIGINAL PAGE IS
OF POOR QUALITYDISTANCE
EC.LON
1994

SEP, ESP
CA, KA
1994

VENUS

1994


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1994

ORIGINAL PAGE 10
OF POOR QUALITY


VENUS 1994



**STA R/S
NON-DSN
1994**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1994



Venus



1995

DECLIN RT.ASC 1995

VENUS

1995

ORIGINAL PAGE IS
OF POOR QUALITY





VENUS

1995

CRISTINA 1995
05/01/01 - 01/16/96

DISTANCE
EC.LON
1995





SEP, ESP
CA, KA
1995

VENUS

1995

OF POOR QUALITY




STA R/S
DSN
1995

VENUS

1995


ORIGINAL PAGE
OF POOR QUALITY



**STA R/S
NON-DSN
1995**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1995



Venus



1996

DECLIN RT.ASC 1996

VENUS

1996

ORIGINAL PAGE 19
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

1996

DISTANCE
EC.LON
1996





SEP, ESP
CA, KA
1996

VENUS

1996


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1996

ORIGINAL SET TO
OF POOR QUALITY


VENUS 1996



**STA R/S
NON-DSN
1996**

ORIGINAL PAGE 13
OF POOR QUALITY

VENUS 1996



Venus



1997

DECLIN RT.ASC 1997

VENUS

1997

ORIGINAL PAGE 19
OF POOR QUALITY





VENUS

1997

ORIGINAL PAGE IS
OF POOR QUALITY

**DISTANCE
EC.LON
1997**





SEP, ESP
CA, KA
1997

VENUS

1997


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
JSN
1997

ORIGINAL PAGE 17
OF POOR QUALITY


VENUS 1997



**STA R/S
NON-DSN
1997**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1997



Venus



1998

DECLIN RT.ASC 1998

/ENUS

1998

ORIGINAL PRINT IS
OF POOR QUALITY





VENUS

1998

**DISTANCE
EC.LON
1998**

ORIGINAL PAGE IS
OF POOR QUALITY





SEP, ESP
CA, KA
1998

VENUS

1998


ORIGINAL PAGE 18
OF POOR QUALITY



STA R/S
DSN
1998

ORIGINAL
OF FOUR


VENUS 1998



STA R/S
NON-DSN
1998

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1998



Venus



1999

DECLIN RT.ASC 1999

VENUS

1999

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IN
OF POOR QUALITY

VENUS

1999

DISTANCE
EC.LON
1999





SEP, ESP
CA, KA
1999

VENUS

1999


ORIGINAL FILE
OF POOR QUALITY



STA R/S
DSN
1999

ORIGINAL COPY
OF POOR QUALITY


VENUS 1999



**STAR/S
NON-DSN
1999**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 1999



Venus



2000

DECLIN RT.ASC 2000

VENUS

2000

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2000

DISTANCE
EC.LON
2000





SEP, ESP
CA, KA
2000

VENUS

2000


ORIGINAL PAGE
OF POOR QUALITY



STA R/S
DSN
2000

ORIGINAL PAGE IS
OF POOR QUALITY


VENUS 2000



**STAR/S
NON-DSN
2000**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2000



Venus



2001

DECLIN RT.ASC 2001

VENUS

2001

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2001

DISTANCE
EC.LON
2001





SEP, ESP
CA, KA
2001

VENUS


2001

ORIGINAL PAGE 13
OF POOR QUALITY



ORIGINAL PAGE 12
OF POOR QUALITY


VENUS 2001



STAR/S
NON-DSN
2001

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2001



Venus



2002

DECLIN RT.ASC 2002

VENUS

2002

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE 13
OF POOR QUALITY

VENUS

2002

DISTANCE
EC.LON
2002





SEP, ESP
CA, KA
2002

VENUS

2002

ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL FILED
OF PCO2


Figure 1 is a line graph showing the variation of station rise/set times (GMT, HR) versus year/month/day (YR/M/D) for four stations: SET 14, RISE 43, SET 63, and RISE 14. The Y-axis represents the time in GMT, HR, ranging from 0 to 24. The X-axis represents the date, ranging from 02/01/01 to 03/01/16. The graph shows seasonal variations in rise and set times, with a shaded region indicating a specific period of interest.

YR/M/D	SET 14 (GMT, HR)	RISE 43 (GMT, HR)	SET 63 (GMT, HR)	RISE 14 (GMT, HR)
02/01/01	16.0	8.0	8.0	8.0
02/01/21	18.0	9.0	8.5	8.5
02/02/10	20.0	9.0	8.0	8.0
02/03/02	22.0	8.5	7.5	7.5
02/03/22	24.0	8.0	7.0	7.0
02/04/11	22.0	8.0	7.0	7.0
02/05/01	20.0	8.5	7.5	7.5
02/05/21	18.0	9.0	8.0	8.0
02/06/10	16.0	10.0	9.0	9.0
02/06/30	14.0	11.0	10.0	10.0
02/07/20	12.0	11.0	10.0	10.0
02/08/09	10.0	10.5	9.5	9.5
02/08/29	8.0	10.0	9.0	9.0
02/09/18	6.0	9.0	8.0	8.0
02/10/08	4.0	7.0	6.0	6.0
02/10/28	2.0	5.0	4.0	4.0
02/11/17	0.0	3.0	2.0	2.0
02/12/07	0.0	1.0	0.0	0.0
02/12/27	0.0	0.0	0.0	0.0
03/01/16	0.0	0.0	0.0	0.0

**STA R/S
NON-DSN
2002**

ORIGINAL PAGE 19
OF POOR QUALITY

VENUS 2002



Venus



2003

DECLIN RT.ASC 2003

VENUS

2003

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE 13
OF POOR QUALITY

VENUS

2003

DISTANCE
EC.LON
2003





SEP, ESP
CA, KA
2003

VENUS

2003


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2003

ORIGINAL PAGE 17
OF PAGE 17


VENUS 2003



STA R/S
NON-DSN
2003

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2003



Venus



2004

DECLIN RT.ASC 2004

VENUS

2004

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE 13
OF POOR QUALITY

VENUS

2004

DISTANCE
EC.LON
2004





SEP, ESP
CA, KA
2004

VENUS


2004

ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY


VENUS 2004



STA R/S
NON-DSN
2004

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2004



Venus



2005

DECLIN
RT.ASC
2005

VENUS

2005

ORIGINAL PAGE 13
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2005

**DISTANCE
EC.LON
2005**





SEP, ESP
CA, KA
2005

VENUS

2005


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2005

ORIGINAL PLAN
OF POOR QUALITY

VENUS 2005




STA R/S
NON-DSN
2005

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2005



Venus



2006



DECLIN
RT.ASC
2006

VENUS

2006

ORIGINAL PAGE 19
OF POOR QUALITY







SEP, ESP
CA, KA
2006

VENUS

2006


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2006

ORIGINAL PAGE 13
OF POOR QUALITY


VENUS 2006



STA R/S
NON-DSN
2006

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2006



Venus



2007

DECLIN
RT.ASC
2007

VENUS

2007

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2007

DISTANCE
EC.LON
2007





SEP, ESP
CA, KA
2007

VENUS


2007

ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY


VENUS 2007



STA R/S
NON-DSN
2007

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2007



Venus



2008

DECLIN RT.ASC 2008

VENUS

2008

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE 19
OF POOR QUALITY

VENUS

2008

DISTANCE
EC.LON
2008





SEP, ESP
CA, KA
2008

VLENUS

2008


ORIGINAL PAGE 10
OF POOR QUALITY



STA R/S
DSN
2008

ORIGINAL PLOTTING
OF POOR QUALITY


VENUS 2008



**STA R/S
NON-DSN
2008**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2008



Venus



2009

DECLIN RT.ASC 2009

VENUS

2009

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL TABLE
OF POOR QUALITY

VENUS

2009

**DISTANCE
EC.LON
2009**





SEP, ESP
CA, KA
2009

ANUS


2009

ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY


VENUS 2009



**STA R/S
NON-DSN
2009**

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2009



Venus



2010

DECLIN RT.ASC 2010

VENUS

2010

ORIGINAL PAGE IS
OF POOR QUALITY





VENUS

2010

**DISTANCE
EC.LON
2010**

ORIGINAL PAGE IS
OF POOR QUALITY





SEP, ESP
CA, KA
2010

VENUS

2010


ORIGINAL PAGE 18
OF POOR QUALITY



STAR/S
DSN
2010

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2010




STA R/S
NON-DSN
2010

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2010



Venus



2011

DECLIN RT.ASC 2011

VENUS

2011

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE 19
OF POOR QUALITY

VENUS

2011

DISTANCE
EC.LON
2011





SEP, ESP
CA, KA
2011

VENUS

2011


ORIGINAL PAGE 19
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STA R/S
DSN
2011


VENUS 2011



STA R/S
NON-DSN
2011

ORIGINAL PAGE 19
OF POOR QUALITY

VENUS 2011



Venus



2012



DECLIN
RT.ASC
2012

VENUS

2012

ORIGINAL PAGE 19
OF POOR QUALITY







SEP, ESP
CA, KA
2012

VENUS

2012


ORIGINAL PAGE 19
OF POOR QUALITY



STA R/S
DSN
2012

ORIGINAL PAGE 1
OF POOR QUALITY


VENUS 2012



STA R/S
NON-DSN
2012

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2012



Venus



2013

DECLIN RT.ASC 2013

VENUS

2013

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2013

**DISTANCE
EC.LON
2013**





SEP, ESP
CA, KA
2013

VENUS

2013


ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

STAR/S
DSN
2013

VENUS 2013




STA R/S
NON-DSN
2013

ORIGINAL PAGE 18
OF POOR QUALITY

VENUS

2013



Venus



2014



DECLIN RT.ASC 2014

VENUS

2014

ORIGINAL PAGE 18
OF POOR QUALITY







SEP, ESP
CA, KA
2014

VENUS

2014

ORIGINAL PAGE IS
OF POOR QUALITY




STA R/S
DSN
2014

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS


2014



**STA R/S
NON-DSN
2014**

ORIGINAL PAGE 18
OF POOR QUALITY

VENUS 2014



Venus



2015

DECLIN RT.ASC 2015

VENUS

2015

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2015

DISTANCE
EC.LON
2015





SEP, ESP
CA, KA
2015

VENUS

2015


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2015

ORIGINAL PAGE 19
OF POOR QUALITY

VENUS 2015




STA R/S
NON-DSN
2015

ORIGINAL PAGE 19
OF POOR QUALITY

VENUS

2015



Venus



2016

DECLIN RT.ASC 2016

VENUS

2016

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE 19
OF POOR QUALITY

VENUS

2016

DISTANCE
EC.LON
2016





SEP, ESP
CA, KA
2016

VENUS

2016

ORIGINAL PAGE IS
OF POOR QUALITY




STA R/S
DSN
2016

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS

2016




STA R/S
NON-DSN
2016

ORIGINAL PAGE 18
OF POOR QUALITY

VENUS

2016



Venus

2017



C-3

DECLIN RT.ASC 2017

VENUS



2017

ORIGINAL PAGE 13
OF POOR QUALITY



VENUS

2017



DISTANCE
EC.LON
2017ORIGINAL PAGE 19
OF POOR QUALITY

SEP, ESP
CA, KA
2017

VENUS


2017

ORIGINAL PAGE 19
OF POOR QUALITY



ORIGINAL PAGE 18
OF POOR QUALITY


VENUS 2017



**STA R/S
NON-DSN
2017**

ORIGINAL PAGE 19
OF POOR QUALITY

VENUS 2017



Venus



2018

DECLIN RT.ASC 2018

VENUS

2018

ORIGINAL PAGE IS
OF POOR QUALITY





VENUS

2018

ORIGINAL PAGE IS
OF POOR QUALITY

**DISTANCE
EC.LON
2018**





SEP, ESP
CA, KA
2018

VENUS

2016


ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STA R/S
DSN
2018


VENUS 2018



STA R/S
NON-DSN
2018

ORIGINAL PAGE 15
OF POOR QUALITY

VENUS 2018



Venus



2019

DECLIN RT.ASC 2019

VENUS

2019

ORIGINAL PAGE IS
OF POOR QUALITY





VENUS

2019

ORIGINAL PAGE IS
OF POOR QUALITY

**DISTANCE
EC.LON
2019**





**SEP, ESP
CA, KA
2019**

VENUS

2019


ORIGINAL PAGE 19
OF POOR QUALITY



STA R/S
DSN
2019

ORIGINAL PAGE IS
OF POOR QUALITY

VENUS 2019




STA R/S
NON-DSN
2019

ORIGINAL PAGE 18
OF POOR QUALITY

VENUS

2019



Venus



2020

DECLIN RT.ASC 2020

VENUS

2020

ORIGINAL PAGE IS
OF POOR QUALITY





VENUS

2020

ORIGINAL PAGE 13
OF POOR QUALITY

**DISTANCE
EC.LON
2020**





SEP, ESP
CA, KA
2020

VENUS

2020


ORIGINAL PAGE 18
OF POOR QUALITY




STA R/S
DSN
2020

ORIGINAL PAGE 13
OF POOR QUALITY

VENUS 2020



VENUS 2020




Positional Data

**Earth
1985—2020**

EC.LON EARTH ALL YEARS

ORIGINAL PAGE 19
OF POOR QUALITY



PRECEDING PAGE BLANK NOT FILMED

PAGE 214 INTENTIONALLY BLANK